Images in
 Cardiovascular Medicine

Clinical and Histological Response to Immunosuppressive Therapy in Giant Cell Myocarditis

You-Jung Choi , MD, and Hyun-Jai Cho , MD, PhD

Division of Cardiology, Department of Internal Medicine, Seoul National University Hospital, Seoul, Korea

OPEN ACCESS

Received: Aug 31, 2018

Revised: Oct 30, 2018

Accepted: Nov 27, 2018

Correspondence to

Hyun-Jai Cho, MD, PhD

Division of Cardiology, Department of Internal Medicine, Seoul National University Hospital, 101, Daehak-ro, Jongno-gu, Seoul 03080, Korea.

E-mail: hyunjaicho@snu.ac.kr
hyunjaicho@gmail.com

A 66-year-old woman was resuscitated from a cardiac arrest after completing 1 cycle of cardiopulmonary resuscitation (**Supplementary Figure 1**). Echocardiography showed severe global akinesia and small pericardial effusion. Coronary angiography was normal. Endomyocardial biopsy was performed. Myocardial pathology confirmed her diagnosis of giant cell myocarditis (GCM) (**Figure 1**). As a treatment for GCM, she received anti-thymocyte globulin (ATG) followed by cyclosporin A and everolimus (**Supplementary Figure 2**).¹⁾ Follow-up pathological examination showed resolving myocarditis on the 12th hospital day (**Figure 2**) at 5 weeks after starting treatment. Histological confirmation is essential for verifying GCM in myocarditis.²⁾ Observations in experimental GCM on Lewis rate model suggest that autoimmune mechanisms mediated by CD4⁺ T lymphocytes are responsible

Figure 1. Histological findings on endomyocardial biopsy. (A) Diffuse geographic myocarditis with severe necrotic change and inflammatory infiltrate (H&E, ×100). (B) Numerous multinucleated giant cells in nongranulomatous background. (C) Collection of inflammatory cells including variable numbers of giant cells (arrows), histiocyte, T-lymphocyte, and eosinophils. (D) The area with necrotic change and severe inflammation showed negative of desmin. (E) Histiocyte infiltration in myocardium (CD68 staining). (F) T-lymphocyte infiltrate in myocardium (CD3, Pan-T cell marker staining). H&E = hematoxylin and eosin.

Copyright © 2019. The Korean Society of Cardiology
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

You-Jung Choi
<https://orcid.org/0000-0003-3512-855X>
Hyun-Jai Cho
<https://orcid.org/0000-0002-2779-4037>

Conflict of Interest

The authors have no financial conflicts of interest.

Author Contributions

Conceptualization: Cho HJ; Data curation: Choi YJ, Cho HJ; Supervision: Cho HJ; Visualization: Choi YJ; Writing - original draft: Choi YJ, Cho HJ; Writing - review & editing: Choi YJ, Cho HJ.

Figure 2. Follow-up endomyocardial biopsy. (A) On 12th day from starting immunosuppressive therapy, myocardial biopsy revealed a marked reduction of the inflammatory infiltrate and the absence of giant cells. The fibrotic and fatty change are seen in the area, where there was necrotic change before treatment (H&E, ×200). (B) On 46th day, massive granulation tissue and fatty change without active myocardial damage was observed (H&E, ×400). H&E = hematoxylin and eosin.

for the pathogenesis of GCM.³⁾ In acute phase, T helper cell type 1 (Th1) is dominant, which may lead to inflammation at the induction of myocarditis. ATG therapy induces a profound decrease in Th1 cell counts.⁴⁾ Early ATG administered seems to play a key role in alleviating the fulminant inflammation of the heart and resulted in a better clinical course.

SUPPLEMENTARY MATERIALS

Supplementary Figure 1

The electrocardiogram (A) and chest x-ray (B) on the first day to the hospital. (A) Sinus rhythm with the low voltage of QRS complexes in the precordial and limb leads. There are no significant ST-segment changes. (B) The chest radiograph shows cardiomegaly with bilateral diffuse infiltrates which suggest cardiogenic pulmonary edema.

[Click here to view](#)

Supplementary Figure 2

Summary for administration of immunosuppressive drugs.

[Click here to view](#)

REFERENCES

1. Fluschnik N, Escher F, Blankenberg S, Westermann D. Fatal recurrence of fulminant giant cell myocarditis and recovery after initialisation of an alternative immunosuppressive regime. *BMJ Case Rep* 2014;2014:bcr2014206386.
2. Vaideeswar P, Cooper LT. Giant cell myocarditis: clinical and pathological features in an Indian population. *Cardiovasc Pathol* 2013;22:70-4.
[PUBMED](#) | [CROSSREF](#)
3. Kodama M, Matsumoto Y, Fujiwara M, Masani F, Izumi T, Shibata A. A novel experimental model of giant cell myocarditis induced in rats by immunization with cardiac myosin fraction. *Clin Immunol Immunopathol* 1990;57:250-62.
[PUBMED](#) | [CROSSREF](#)

4. Weimer R, Staak A, Süsal C, et al. ATG induction therapy: long-term effects on Th1 but not on Th2 responses. *Transpl Int* 2005;18:226-36.

[PUBMED](#) | [CROSSREF](#)