Erratum

Erratum: Significance of Differential Characteristics in Infantile Kawasaki Disease

Ji Hee Kwak , MD¹, JungHwa Lee , MD, PhD², and Kee Soo Ha , MD, PhD²

¹Department of Pediatrics, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, Seoul, Korea

²Department of Pediatrics, Korea University Guro Hospital, Korea University College of Medicine, Seoul, Korea

OPEN ACCESS

► This corrects the article “Significance of Differential Characteristics in Infantile Kawasaki Disease” in volume 49 on page 755.

Copyright © 2020. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Ji Hee Kwak

<https://orcid.org/0000-0001-6376-0427>

JungHwa Lee

<https://orcid.org/0000-0002-6592-3653>

Kee Soo Ha

<https://orcid.org/0000-0001-6753-5411>

In the article, “Significance of differential characteristics in infantile Kawasaki disease” in volume 49(8), page 755-765,

In the process of inputting standard deviation (SD) in **Table 1**, each 1,000 omitted. The units were entered incorrectly without checking the footnote of Table A1.20 in the reference article 9. The true Z value = (measured value – mean value) / original incorrect SD × 1,000 and therefore, the Z-values should be changed to 1/1,000 of the original incorrect SD, respectively in Tables 6 and 7. The original p values do not need to correct since all groups were compared with simultaneously 1,000 multiplied values.

Before correction

We deeply apologize for any inconvenience it may have caused.

Table 1. Means and SDs of white cell counts with differential and red cell counts according to ages

Ages	White cells (/μL)					Ages	Sex	Red cells (g/dL)
	Total leukocytes	Neutrophils	Lymphocytes	Monocytes	Eosinophils			Hemoglobins
	Mean±SD	Mean±SD	Mean±SD	Mean±SD	Mean±SD			Mean±SD
1–3 months	10,800±3.63	3,800±2.00	6,000±3.50	700±0.23	300±0.2	1–2 months	-	14±2
4–8 months	11,900±2.88	3,800±1.88	7,300±2.34	600±0.23	300±0.2	3–6 months	-	11.5±1
9–17 months	11,400±2.88	3,500±1.75	7,000±1.63	600±0.28	300±0.14	6–59 months	-	12.5±0.75
18–35 months	10,600±2.75	3,500±1.75	6,300±1.63	500±0.20	300±0.14	60–95 months	-	13±0.75
36–59 months	9,100±2.50	3,800±1.75	4,500±1.50	500±0.20	300±0.23	96–143 months	-	13.5±0.75
60–83 months	8,500±2.34	4,300±1.63	3,500±1.38	400±0.19	200±0.21	12–14 years	M	14±0.75
84–107 months	8,300±2.25	4,400±1.63	3,300±1.33	400±0.21	200±0.25	12–14 years	F	13.5±0.75
108–156 months	8,100±2.25	4,400±1.55	3,100±1.25	400±0.22	200±0.24	15–17 years	M	15±1
157–221 months	7,800±2.13	4,400±1.55	2,800±1.00	400±0.14	200±0.12	15–17 years	F	14±1
≥222 months	7,400±1.63	4,400±1.48	2,500±0.95	400±0.17	200±0.07	-	-	-

Table 6. Usefulness and comparison of Z in infants and non-infants with KD compared with FC groups in pre-IVIG phase

Variables	Infants (n=216)			Non-infants (n=757)			FC infants vs. FC non-infants
	KD (n=192)	FC (n=24)	p value*	KD (n=667)	FC (n=90)	p value*	p value†
Z-leukocyte	815 (–174, 2,060)	–734 (–2,257, 1,710)	0.008	815 (–174, 2,060)	1,360 (182, 2,800)	<0.001	0.640
Z-neutrophil	2,597 (1,135, 3,967)	224 (–1,201, 1,944)	<0.001	2,597 (1,135, 3,967)	1,360 (182, 2,800)	<0.001	0.400
Z-lymphocyte	–1,309 (–1,986, –497)	–1,623 (–2,498, 76)	0.700	–1,309 (–1,986, –497)	–1,718 (–2,263, –1,077)	0.290	0.740
Z-eosinophil	76 (–896, 1,581)	–999 (–1,404, 736)	0.040	76 (–896, 1,581)	–520 (–1,193, 750)	<0.001	0.340
Z-monocyte	854 (–336, 2,415)	1,319 (282, 2,574)	0.390	854 (–336, 2,415)	688 (–442, 2,054)	0.080	0.780
NLR	1.92 (1.27, 3.06)	0.91 (0.41, 1.31)	<0.001	1.92 (1.27, 3.06)	3.43 (2.10, 5.98)	<0.001	0.002
NER	22 (12, 55)	17 (10, 49)	0.370	22 (12, 55)	41 (19, 119)	0.030	0.004
R-hemoglobin	10.9 (10.3, 11.5)	11.6 (10.4, 11.9)	0.060	10.9 (10.3, 11.5)	11.5 (10.9, 12.1)	<0.001	0.020
Z-hemoglobin	–1.5 (–2.7, –0.8)	–0.0 (–0.1, 0.0)	<0.001	–1.5 (–2.7, –0.8)	–1.5 (–2.2, –0.7)	<0.001	0.740
PLT	384 (314, 466)	304 (185, 388)	0.001	384 (314, 466)	328 (275, 395)	<0.001	0.210
CRP	76 (36, 123)	17 (2, 67)	<0.001	76 (36, 123)	79 (47, 128)	<0.001	0.170
ESR	66 (46, 94)	34 (11, 72)	0.004	66 (46, 94)	90 (57, 100)	<0.001	0.350

Table 7. Laboratory characteristics of KD infants compared with KD non-infants in pre-IVIG, post-IVIG, and chronic phases

Variables	Pre-IVIG phase			Post-IVIG phase			Chronic phase		
	Infants (n=192)	Non-infants (n=667)	p value*	Infants (n=192)	Non-infants (n=667)	p value*	Infants (n=192)	Non-infants (n=667)	p value*
Z-leukocyte	816 (–174, 2,060)	1,360 (189, 2,800)	0.002	–976 (–1,783, –177)	–916 (–1,496, –116)	0.150	–944 (–1,394, –429)	–680 (–1,236, –182)	0.020
Z-neutrophil	2,598 (1,135, 3,967)	3,217 (1,659, 5,301)	<0.001	–946 (–1,419, –220)	–599 (–1,169, 303)	0.001	–909 (–1,195, –567)	–469 (–938, 117)	<0.001
Z-lymphocyte	–1,309 (–1,986, –497)	–1,715 (–2,259, –1,075)	<0.001	–1,060 (–1,690, –338)	–1,142 (–1,736, –563)	0.230	–829 (–1,318, –289)	–739 (–1,422, –140)	0.710
Z-eosinophil	76 (–897, 1,581)	–520 (–1,182, 765)	<0.001	861 (–174, 2,032)	434 (–378, 1,368)	0.005	492 (–197, 1,747)	–199 (–680, 653)	<0.001
Z-monocyte	854 (–336, 2,415)	688 (–442, 2,057)	0.260	404 (–607, 1,624)	377 (–406, 1,287)	0.740	21 (–663, 531)	133 (–437, 815)	0.220
NLR	1.92 (1.27, 3.06)	3.43 (2.10, 5.98)	<0.001	0.4 (0.2, 0.7)	0.8 (0.5, 1.3)	<0.001	0.4 (0.3, 0.5)	0.7 (0.5, 1.0)	<0.001
NER	22 (12, 58)	41 (19, 119)	<0.001	3.9 (2.2, 8.8)	6.9 (4.2, 13.8)	<0.001	4.8 (2.8, 8.0)	10.3 (6.6, 16.9)	<0.001
R-hemoglobin	11 (10.3, 11.5)	11.5 (10.9, 12.1)	<0.001	10.2 (9.7, 10.9)	11.0 (10.4, 11.6)	<0.001	11.1 (10.6, 11.6)	11.6 (11.2, 12.2)	<0.001
Z-hemoglobin	–1.7 (–2.7, –0.8)	–1.5 (–2.2, –1.0)	0.040	–2.5 (–3.5, –1.6)	–2.1 (–2.9, –1.4)	0.003	–1.6 (–2.5, –0.9)	–1.3 (–1.9, –0.5)	0.006
PLT	384 (314, 466)	328 (275, 395)	<0.001	505 (416, 638)	461 (358, 552)	<0.001	380 (299, 519)	330 (277, 407)	0.001
CRP	76 (36, 123)	79 (47, 128)	0.200	18 (8, 32)	17 (8, 34)	0.640	0.3 (0.2, 1.2)	0.5 (0.3, 2.1)	0.004
ESR	66 (46, 94)	90 (57, 100)	<0.001	70 (48, 98)	88 (65, 112)	<0.001	13 (7, 25)	24 (14, 42)	<0.001

After correction

Table 1. Means and SDs of white cell counts with differential and red cell counts according to ages

Ages	White cells (/μL)					Ages	Sex	Red cells (g/dL)
	Total leukocytes	Neutrophils	Lymphocytes	Monocytes	Eosinophils			Hemoglobins
	Mean±SD	Mean±SD	Mean±SD	Mean±SD	Mean±SD			Mean±SD
1–3 months	10,800± 3,630	3,800± 2,000	6,000± 3,500	700± 230	300± 200	1–2 months	-	14±2
4–8 months	11,900± 2,880	3,800± 1,880	7,300± 2,340	600± 230	300± 200	3–6 months	-	11.5±1
9–17 months	11,400± 2,880	3,500± 1,750	7,000± 1,630	600± 280	300± 140	6–59 months	-	12.5±0.75
18–35 months	10,600± 2,750	3,500± 1,750	6,300± 1,630	500± 200	300± 140	60–95 months	-	13±0.75
36–59 months	9,100± 2,500	3,800± 1,750	4,500± 1,500	500± 200	300± 230	96–143 months	-	13.5±0.75
60–83 months	8,500± 2,340	4,300± 1,630	3,500± 1,380	400± 190	200± 210	12–14 years	M	14±0.75
84–107 months	8,300± 2,250	4,400± 1,630	3,300± 1,330	400± 210	200± 250	12–14 years	F	13.5±0.75
108–156 months	8,100± 2,250	4,400± 1,550	3,100± 1,250	400± 220	200± 240	15–17 years	M	15±1
157–221 months	7,800± 2,130	4,400± 1,550	2,800± 1,000	400± 140	200± 120	15–17 years	F	14±1
≥222 months	7,400± 1,630	4,400± 1,480	2,500± 950	400± 170	200± 70	-	-	-

Table 6. Usefulness and comparison of Z in infants and non-infants with KD compared with FC groups in pre-IVIG phase

Variables	Infants (n=216)			Non-infants (n=757)			FC infants vs. FC non-infants
	KD (n=192)	FC (n=24)	p value*	KD (n=667)	FC (n=90)	p value*	p value†
Z-leukocyte	0.8 (–0.2, 2.1)	–0.7 (–2.3, 1.7)	0.008	0.8 (–0.2, 2.1)	1.4 (0.2, 2.8)	<0.001	0.640
Z-neutrophil	2.6 (1.1, 4.0)	0.2 (–1.2, 1.9)	<0.001	3.0 (1.1, 4.0)	1.4 (0.2, 2.8)	<0.001	0.400
Z-lymphocyte	–1.3 (–2.0, –0.5)	–1.6 (–3.0, 0.1)	0.700	–1.31 (–2.0, –0.5)	–1.7 (–2.3, –1.1)	0.290	0.740
Z-eosinophil	0.1 (–0.9, 1.6)	–1.0 (–1.4, 0.7)	0.040	0.1 (–0.9, 1.6)	–0.5 (–1.2, 0.8)	<0.001	0.340
Z-monocyte	0.9 (–0.3, 2.4)	1.3 (0.3, 2.6)	0.390	0.9 (–0.3, 2.4)	0.7 (–0.4, 2.1)	0.080	0.780
NLR	1.9 (1.3, 3.1)	0.9 (0.4, 1.3)	<0.001	1.9 (1.3, 3.1)	3.4 (2.1, 6.0)	<0.001	0.002
NER	22 (12, 55)	17 (10, 49)	0.370	22 (12, 55)	41 (19, 119)	0.030	0.004
R-hemoglobin	10.9 (10.3, 11.5)	11.6 (10.4, 11.9)	0.060	10.9 (10.3, 11.5)	11.5 (10.9, 12.1)	<0.001	0.020
Z-hemoglobin	–1.5 (–2.7, –0.8)	–0.0 (–0.1, 0.0)	<0.001	–1.5 (–2.7, –0.8)	–1.5 (–2.2, –0.7)	<0.001	0.740
PLT	384 (314, 466)	304 (185, 388)	0.001	384 (314, 466)	328 (275, 395)	<0.001	0.210
CRP	76 (36, 123)	17 (2, 67)	<0.001	76 (36, 123)	79 (47, 128)	<0.001	0.170
ESR	66 (46, 94)	34 (11, 72)	0.004	66 (46, 94)	90 (57, 100)	<0.001	0.350

Table 7. Laboratory characteristics of KD infants compared with KD non-infants in pre-IVIG, post-IVIG, and chronic phases

Variables	Pre-IVIG phase			Post-IVIG phase			Chronic phase		
	Infants (n=192)	Non-infants (n=667)	p value*	Infants (n=192)	Non-infants (n=667)	p value*	Infants (n=192)	Non-infants (n=667)	p value*
Z-leukocyte	0.8 (–0.2, 2.1)	1.4 (0.2, 2.8)	0.002	–1.0 (–1.8, –0.2)	–0.9 (–1.5, –0.1)	0.150	–0.9 (–1.4, –0.4)	–0.7 (–1.2, –0.2)	0.020
Z-neutrophil	2.6 (1.1, 4.0)	3.2 (1.7, 5.3)	<0.001	–1.0 (–1.4, –0.2)	–0.6 (–1.2, 0.3)	0.001	–0.9 (–1.2, –0.6)	–0.5 (–0.9, 0.1)	<0.001
Z-lymphocyte	–1.3 (–2.0, –0.5)	–1.7 (–2.3, –1.1)	<0.001	–1.1 (–1.7, –0.3)	–1.1 (–1.7, –0.6)	0.230	–0.8 (–1.3, –0.3)	–0.7 (–1.4, –0.1)	0.710
Z-eosinophil	0.1 (–0.9, 1.6)	–0.5 (–1.2, 0.8)	<0.001	0.9 (–0.2, 2.0)	0.4 (–0.4, 1.4)	0.005	0.5 (–0.2, 1.7)	–0.2 (–0.7, 0.7)	<0.001
Z-monocyte	0.9 (–0.3, 2.4)	0.7 (–0.4, 2.1)	0.260	0.4 (–0.6, 1.6)	0.4 (–0.4, 1.3)	0.740	0.0 (–0.7, 0.5)	0.1 (–0.4, 0.8)	0.220
NLR	1.9 (1.3, 3.1)	3.4 (2.1, 6.0)	<0.001	0.4 (0.2, 0.7)	0.8 (0.5, 1.3)	<0.001	0.4 (0.3, 0.5)	0.7 (0.5, 1.0)	<0.001
NER	22 (12, 58)	41 (19, 119)	<0.001	3.9 (2.2, 8.8)	6.9 (4.2, 13.8)	<0.001	4.8 (2.8, 8.0)	10.3 (6.6, 16.9)	<0.001
R-hemoglobin	11 (10.3, 11.5)	11.5 (10.9, 12.1)	<0.001	10.2 (9.7, 10.9)	11.0 (10.4, 11.6)	<0.001	11.1 (10.6, 11.6)	11.6 (11.2, 12.2)	<0.001
Z-hemoglobin	–1.7 (–2.7, –0.8)	–1.5 (–2.2, –1.0)	0.040	–2.5 (–3.5, –1.6)	–2.1 (–2.9, –1.4)	0.003	–1.6 (–2.5, –0.9)	–1.3 (–1.9, –0.5)	0.006
PLT	384 (314, 466)	328 (275, 395)	<0.001	505 (416, 638)	461 (358, 552)	<0.001	380 (299, 519)	330 (277, 407)	0.001
CRP	76 (36, 123)	79 (47, 128)	0.200	18 (8, 32)	17 (8, 34)	0.640	0.3 (0.2, 1.2)	0.5 (0.3, 2.1)	0.004
ESR	66 (46, 94)	90 (57, 100)	<0.001	70 (48, 98)	88 (65, 112)	<0.001	13 (7, 25)	24 (14, 42)	<0.001