Images in Cardiovascular Medicine

Weeping Heart: Regrowing Thrombus at Basal Left Ventricle in Acute Stroke

Soon Myung Jung , MD¹, Yong Hyun Park , MD, PhD¹, Sang Hyun Lee , MD, PhD¹,
Soo Yong Lee , MD, PhD¹, Kyung-Pil Park , MD, PhD², and June Hong Kim, MD, PhD¹

¹Division of Cardiology, Department of Internal Medicine, Pusan National University Yangsan Hospital, Yangsan, Korea

²Department of Neurology, Pusan National University Yangsan Hospital, Yangsan, Korea

OPEN ACCESS

Received: Jan 13, 2020

Accepted: Jan 22, 2020

Correspondence to

Yong Hyun Park, MD, PhD

Division of Cardiology, Department of Internal Medicine, Pusan National University Yangsan Hospital, 20 Geumo-ro, Yangsan 50612, Korea.
E-mail: nadroj70@gmail.com

Copyright © 2020. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Soon Myung Jung
<https://orcid.org/0000-0003-0846-3133>

Yong Hyun Park
<https://orcid.org/0000-0001-6122-214X>

Sang Hyun Lee
<https://orcid.org/0000-0001-7196-2643>

Soo Yong Lee
<https://orcid.org/0000-0003-2616-1294>

Kyung-Pil Park
<https://orcid.org/0000-0003-4952-3796>

Conflict of Interest

The authors have no financial conflicts of interest.

A 39-year-old male visited the emergency department 90 minutes after the sudden onset of right-sided weakness and aphasia following endoscopic examination for epigastric discomfort and dyspepsia. He had suffered congestive heart failure for years, but was lost to follow-up. Brain magnetic resonance imaging revealed acute embolic infarction at the right cerebrum and left cerebellum. Chest X-ray showed cardiomegaly. The electrocardiogram revealed sinus rhythm and left ventricular (LV) hypertrophy. Laboratory findings were non-specific. On transthoracic echocardiography (TTE), global hypo-kinesis and severe LV systolic dysfunction were found, but specific embolic sources were not evident (**Figure 1A, Supplementary Video 1**). After 3 days of medication with antiplatelet agents, transesophageal echocardiography found a mobile echogenic mass in the basal anterior wall of LV, which was not observed in the previous TTE (**Figure 1B, Supplementary Video 2**). Repeated TTE also found newly appeared mobile echogenic mass at the basal anterior wall of LV (**Figure 1C, Supplementary Video 3**). After a resolution of thrombus with 1 week of adequate anticoagulation therapy, TTE revealed a recess in the basal left ventricle, which had played a role of the nest for a nidus formation (**Figure 1D, Supplementary Video 4**). After 6 days of uneventful recovery, the patient was discharged with warfarin and antihypertensive therapy in good condition. In this case, the combination of decreased LV systolic function and a ventricular recess might be a necessary condition for thrombus formation. Thorough vigilance should be exerted throughout the LV from apex to base to find out the source of cardiac embolism.

SUPPLEMENTARY MATERIALS

Supplementary Video 1

TTE demonstrates no visible embolic source in the LV.

[Click here to view](#)

Supplementary Video 2

TEE showed newly appeared echogenic mobile mass on LV basal anterior wall.

[Click here to view](#)

Author Contributions

Conceptualization: Park YH, Lee SY, Park KP, Kim JH; Supervision: Park YH, Park KP, Kim JH; Writing - original draft: Jung SM; Writing - review & editing: Park YH, Lee SH, Lee SY, Kim JH.

Figure 1. Echocardiographic images. (A) TTE demonstrates no visible embolic source in the LV. (B) TEE showed newly appeared echogenic mobile mass on LV basal anterior wall. (C) TTE also showed newly appeared echogenic mobile mass not seen in previous TTE. (D) After 1 week of anticoagulation, TTE revealed a recess in the basal LV. LV = left ventricle; TEE = transthoracic echocardiography; TTE = transthoracic echocardiography.

Supplementary Video 3

TTE also showed newly appeared echogenic mobile mass not seen in previous TTE.

[Click here to view](#)

Supplementary Video 4

After 1 week of anticoagulation, TTE revealed a recess in the basal LV.

[Click here to view](#)